



SCAT3 Symptom Evaluation

(Athletes ages 13 and older)

Patient Name: _____

DOB: _____

How do you feel?

"You should score yourself on the following symptoms, based on how you feel now".

	None	Mild		Moderate		Severe	
Headache	0	1	2	3	4	5	6
“Pressure in head”	0	1	2	3	4	5	6
Neck Pain	0	1	2	3	4	5	6
Nausea or vomiting	0	1	2	3	4	5	6
Dizziness	0	1	2	3	4	5	6
Blurred vision	0	1	2	3	4	5	6
Balance problems	0	1	2	3	4	5	6
Sensitivity to light	0	1	2	3	4	5	6
Sensitivity to noise	0	1	2	3	4	5	6
Feeling slowed down	0	1	2	3	4	5	6
Feeling like “in a fog”	0	1	2	3	4	5	6
“Don’t feel right”	0	1	2	3	4	5	6
Difficulty concentrating	0	1	2	3	4	5	6
Difficulty remembering	0	1	2	3	4	5	6
Fatigue or low energy	0	1	2	3	4	5	6
Confusion	0	1	2	3	4	5	6
Drowsiness	0	1	2	3	4	5	6
Trouble falling asleep	0	1	2	3	4	5	6
More emotional	0	1	2	3	4	5	6
Irritability	0	1	2	3	4	5	6
Sadness	0	1	2	3	4	5	6
Nervous or Anxious	0	1	2	3	4	5	6

Total number of symptoms (Maximum possible 22)

Symptom severity score (Maximum possible 132)

Do the symptoms get worse with physical activity? ☐ Y ☐ N Do the symptoms get worse with mental activity? ☐ Y ☐ N

☐ self rated

☐ self rated and clinician monitored

☐ clinician interview

☐ self rated with parent input

Overall rating: If you know the athlete well prior to the injury, how different is the athlete acting compared to his/her usual self?

Please select one response:

☐ no different

☐ very different

☐ unsure

☐ N/A

HW-21
REV-9/14

Scoring on the SCAT3 should not be used as a stand-alone method to diagnose concussion, measure recovery or make decisions about an athlete's readiness to return to competition after concussion. Since signs and symptoms may evolve over time, it is important to consider repeat evaluation in the acute assessment of concussion. SCAT3 Sport Concussion Assessment Tool 3.